LIPIDS FOR

INJECTABLES

IOI OLEOCHEMICAL

IOI Oleo GmbH
PHARMA

Injectables – IV, SC and IM

C8/C10 triglycerides (MCT oils) and propylene glycol ester, used as a carrier:

MIGLYOL® 810 N
Medium Chain Triglycerides (ratio ~ 70:30%)

MIGLYOL® 812 N
Medium Chain Triglycerides (ratio ~ 60:40%)

MIGLYOL® 840
Propylene Glycol Dicaprylocaprate

Application examples

- Narcotics (Propofol)
- Malaria (Artemether, Lumefantrin)
- Muscle relaxant (Etomidate)
- API-loaded microspheres (Exenatide)
- Veterinary injection suspension (Amoxicillin, Ceftiofur)

Injectables – Enteral and parenteral nutrition

C8/C10 triglycerides (MCT oils)
MIGLYOL® 812 N –
Medium Chain Triglycerides (ratio ~ 60:40%)

Dosage forms:
enteral and parenteral nutrition

MIGLYOL® 812 N has already been used in enteral and parenteral nutrition for decades.

From a regulatory point of view, in some of these formulations specific ingredients incl. **MIGLYOL® 812 N** are classified as API.

IOI is the holder of the first CEP (Certificate of Suitability) ever granted for a Medium Chain Triglyceride and has filed a US-DMF Type II.

IOI Oleo GmbH
Hamburg, Germany | +49 40 28 00 31-0 | pharma@ioioleo.de | www.ioioleo.de